

MATH 251 (Fall 2004) Exam 3, Nov 29th

No calculators, books or notes! Show all work and give **complete explanations** for all your answers. This 65 minute exam is worth 75 points.

(1) [15 pts]

(a) Suppose that (0, 2) is a critical point of a function g with continuous second partial derivatives. What can you say about g if

 $g_{xx}(0,2) = -1$ $g_{xy}(0,2) = 2$ $g_{yy}(0,2) = -8?$

(b) Find the maximum rate of change of the function $f(x, y, z) = x^2 y^3 z^4$ at the point (1, 1, 1), and the direction in which it occurs.

(2) [15 pts]

(a) Evaluate the line integral $\int_C \mathbf{F} \cdot d\mathbf{r}$ where $\mathbf{F}(x, y) = x^2 y^3 \mathbf{i} - y \sqrt{x} \mathbf{j}$, and where C is the curve parametrized by $\mathbf{r}(t) = t^2 \mathbf{i} - t^3 \mathbf{j}$ for $0 \le t \le 1$.

(b) Use Green's Theorem to evaluate $\int_C \sqrt{1+x^3} \, dx + 2xy \, dy$, where C is the curve that consists of straight lines joining (0,0) to (1,0), (1,0) to (1,3), and (1,3) to (0,0).

(3) [15 pts]

Find the absolute maximum and minimum of the function f(x, y) = xy on the region $3x^2 + 3y^2 \le 1$.

(4) [15 pts] Calculate the integral $\iint_D y \, dA$, where D is the region in the first quadrant that lies above the hyperbola xy = 1, above the line y = x, and below the line y = 2.

(5) [15 pts] Let \mathbf{F} be the vector field

$$\mathbf{F}(x,y) = (2x\cos y - y\cos x)\mathbf{i} + (-x^2\sin y - \sin x)\mathbf{j}.$$

(a) Determine whether or not **F** is conservative. If it is, find a function f so that $\mathbf{F} = \nabla f$.

(b) Let C be the curve that is the straight line from (0,0) to (1,1). What is $\int_C \mathbf{F} \cdot d\mathbf{r}$?

Pledge: I have neither given nor received aid on this exam

Signature: _____